



# Examination of the Physical Aspects of Hydrogen Storage in MOFs

Omar M. Yaghi and Adam J. Matzger  
Department of Chemistry  
University of Michigan  
Ann Arbor, MI 48109

This presentation does not contain any proprietary or confidential information

Project ID #  
STP39 Yaghi



# Overview

## Timeline

- Project start date  
1/1/2005
- Project end date  
12/31/2009
- Percent complete  
2.5%

## Budget

- Total project funding
  - DOE share: \$ 1.75M
  - Contractor share: \$ 0.437M
- Funding received in FY04
  - \$0.00
- Funding for FY05
  - \$62,500

## Barriers

- Technical barriers addressed
  - B) Weight and Volume
  - C) Efficiency
  - M) Hydrogen Capacity and Reversibility
  - N) Lack of Understanding of Hydrogen Physisorption and Chemisorption
- Technical targets by YR 2010
  - Gravimetric capacity: 6.0%
  - Volumetric capacity: 4.5%
  - Operating ambient temp.: -30/50 °C
  - Cycle life: 1000


## Partners

- NREL Team
- Yue Wu (University of North Carolina)
- Hansong Chen (Air Products)



# CbHS Center of Excellence Partners

*9 university projects (at 7 universities), 4 government labs, 1 industrial partner*





# Objectives

***To develop novel, highly porous metal-organic framework materials (MOFs) as high capacity sorbents for H<sub>2</sub> storage applications.***

- Assess gravimetric and volumetric H<sub>2</sub> storage capacities of MOFs.
  - Explore pressure and temperature dependence of H<sub>2</sub> uptake in existing MOF materials over the parameter range specified in DOE YR2010 guidelines.
- Determine the optimal pore size and functionality for H<sub>2</sub> sorption and release in MOFs.
  - Characterize H<sub>2</sub> adsorption sites in existing MOF materials.
  - Study the relationship between pore size and level of H<sub>2</sub> uptake.



# Approach



- Equilibrium  $H_2$  uptake as a function of structure.
  - Survey hydrogen uptake levels in existing MOFs, under a variety of conditions, to determine promising materials.
  - Use Raman Spectroscopy to probe  $H_2$  sorbed in pores.
- Correlate systematic changes in organic links with uptake.
  - Use above results to aid design of new structures with greater hydrogen storage capacities.
  - Synthesize new organic linkers and the corresponding MOFs.
  - Measure equilibrium  $H_2$  uptake.



# General Synthetic Strategy of MOFs





Inorganic Units






SBU



Organic Units




SBU






# Control of MOF Dimensionality


1D



2D




3D










# Synthesis of an Isoreticular Series of MOFs







# Variation in MOF Metrical Parameters






# Mechanism of H<sub>2</sub> Uptake in MOFs

## Chemisorption

Mg-Hydride, 300 °C


Weight % H<sub>2</sub>



## Physisorption


MOF-5, 24 °C

Weight % H<sub>2</sub>







# Inelastic Neutron Scattering of $H_2$ in MOF-5



*Smallest repeat unit of MOF-5*




*Extended structure of MOF-5*








# H<sub>2</sub> Sorption at 1 atm & 77 K







# Synthesis of a Cu-based MOF







# MOF-505 H<sub>2</sub> (77 K) Sorption Isotherms– Activation Study





*MOF-505*





# Future Work

TASK	2005	2006	2007	2008	2009
<b>Task 1: Equilibrium <math>H_2</math> Uptake vs Structure</b> Measure temperature and pressure dependence of $H_2$ uptake in 4 existing MOFs Computational design of new linkers Explore new linkers		 Go/ No-go Option	 Go/ No-go Option		
<b>Task 2: Thermodynamics &amp; Kinetics <math>H_2</math> Uptake</b> Determine $H_2$ binding energy to framework by sorption Analysis $H_2$ uptake rates in single crystals by Raman spectroscopy Measure $H_2$ flux in single x-tals					
<b>Task 3: Mechanism of <math>H_2</math> Uptake</b> Analysis $H_2$ binding sites in MOFs by Raman spectroscopy NMR measurement of $H_2$ uptake					
<b>Task 4: Down-select Optimum Materials</b> Optimize scale-up synthesis and activation Cost analysis of material					



# Publications and Presentations

---

Please list any publications and presentations that have resulted from work on this project.

- *No publications resulting from current funding at this stage of the project.*



# Hydrogen Safety

*The most significant hydrogen hazard associated with this project is:*

- High exposure to  $H_2$  gas with possibility of personal injury due to decreased oxygen content in the atmosphere.
- High concentrations of  $H_2$  may pose a fire or explosion in and around instrumentation.



# Hydrogen Safety

*Our approach to deal with this hazard:*

- Dedicated a single laboratory for all H<sub>2</sub> experiments.
- Installed active ventilation snorkles from laboratory hoods to all instrumentation consuming/ releasing H<sub>2</sub>.
- Installed atmospheric H<sub>2</sub> detector (ppm level detection) outfitted with an alarm in the dedicated laboratory.



# Acknowledgements

